National Aeronautics and Space Administration Goddard Space Flight Center Contract No.NAS-5-3760

ST - CC - 10 390

NASA TT F-9902

•	N65-33824				
FORM &C	(ACCESSION NUMBER)	(THRU)			
AGILITY	(PAGES)	24			
-	(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)			

MEASUREMENT BY Y — SPECTROMETRY OF THE RADIOACTIVITY OF THE GRANES METEORITE FALLEN ON 13 NOVEMBER 1964

by
Daniel Nordemann
Jâcques Tobailem
Michele Schineizer

[FRANCE]

GPO PRICE \$	
CSFTI PRICE(S) \$	
Hard copy (HC) Microfiche (MF) ff 653 July 65	1.00

24 SEPTEMBER 1965

MEASUREMENT BY Y — SPECTROMETRY OF THE RADIOACTIVITY OF THE GRANES METEORITE FALLEN ON 13 NOVEMBER 1964 *

Comptes-Rendus de l'Académie des Sciences NETEORITES, T. 260, pp. 6664-6, Paris, 21 June 1965 by Daniel Nordemann
Jacques Tobailem
Michèle Schineizer

SUMMARY

33824

The activities of the Y - emitters, present in a 1.186 kg sample of the Granès (Aude) stone meteorite, have been determined by quantitative Y - spectrometry of weak proper motion.

On 13 November 1964, at about 1700 hours, a stone meteorite (chondrite) has fallen in the village of Granes, near Quillan (Aude). The total mass, recovered in several fragments, was of the order of 9 kg.


Thanks to M. Rémy of the Faculty of Sciences of Montpellier, a fragment of 1.186 kg could reach our laboratory only seven days after the fall. Nondesctructive γ - spectrometry measurements with high sensitivity were immediately undertaken.

Such measurements have already allowed us to detect and determine the principal γ — emitting radioelements produced in the meteorites by action of cosmic radiation [1, 2].

The sample has been measured during four months by 10-hour intervals so as to ascertain the decay of short-lived radioactive elements. The figure, next page, shows one of the first spectrograms obtained.

^{*} La radioactivité de la météorite Granès (chute du 13 novembre 1964) mesurée par spectrométrie Y.

The presence of aluminum 26, sodium 22, manganese 54, scandium 46 and potassium 40 is clearly ascertained by their characteristic photoelectric peaks.



Y - Spectrum of the Granes Meteorite.

(Fall of 13 November 1964, 70-hour measurement, having begun on 20 November 1964 on a sample weighing 1.186 kg, placed on a scintillation crystal NaI (Ti) of 10×8 cm dimensions; adjustment: 15 keV/channel.

Other emitters, of shorter periods, manganese 52, vanadium 48, chromium 51, cobalt 58, cobalt 56 are detected by the analysis of counting rate decrease in the different energy ranges. This procedure allows to separate in the photoelectric peaks the contributions of the emitters, whose periods are sufficiently short to offer an appreciable decrease in the course of measurement duration.

Owing to the calibration of the installation in the measurement conditions, these determinations are quantitative. —
This calibration was effected by means of calibrated sources of mercury 203, manganese 54, cobalt 60 and potassium 40, distributed in iron powder dummies simulating the shape and the electron density of the sample.

The results of the aggregate of measurements are compiled in the following table:

EMITTER		ACTIVITIES (dpm.kg-1).	PRINCIPAL MODES OF FORM			
⁵² Mn	5,6 jours days	6 ±6	⁵⁶ Fe(p, 2p 3n) ⁵² Mn; ⁵⁶ Fe(p, 2pn) ⁵² Mn			
48V	16,1 »	17 ±6	$^{56}\text{Fe}(p,4p5n)$ ⁴⁸ V			
51 Cr	27,8 »	79 ± 25	56 Fe $(p, 3p 3n)$ 51 Cr			
³⁸ Со	71,3 » 73,3 »	20 ± 5	$\begin{cases} {}^{58}\text{Ni}(n,p) {}^{58}\text{Co} \\ {}^{56}\text{Fe}(p,n) {}^{56}\text{Co} \end{cases}$			
46Sc	84,0 »	$5,5\pm 2$	⁵⁵ Fe(p, 6p 5n) ⁴⁶ Sc			
54Mn	313,5 »	50 ±7	54 Fe (n, p) 54 Mn;			
²² Na	2,58 ans 7,38.10 ⁵ »	± 10	56 Fe $(n, p 2n)$ 54 Mn 28 Si $(p, 4p 3n)$ 22 Na			
²⁶ A1	7,38.10 ⁵ »	81 ± 10	²⁸ Si(p, 2pn) ²⁶ Al			
⁴ºK	1,27.10° »	1565 ± 75	Nucléosynthèse; [nucleosynthesis] 56 Fe $(p, 8p9n)$ 16 K; 10 Ca (p, n) 10 K			

The radioelement ⁵²Mn, measured in the Granes meteorite in the course of the present work, had not yet been detected in the meteorites; this is due to its short period (5 - 6 days) and to the delays, separating the analyses from the fall. Its rate of activity can provide information on the recent past of meteorite's exposure to cosmic radiation.

The effective formation sections of manganese 52 and manganese 54 starting from iron have been determined in different irradiation conditions. Their ratio, as a function of proton energy, is: 0.39 at 730 MeV, according to Honda and Lal [3], and 0.50; 0.30; 0.37 and 0.37, respectively at 800, 1500, 2200 and 2900 MeV after Rayudu [4]. These values imply that one should have been expecting to find an activity in manganese 52 at least equal to about one third of that of manganese 54, and this in case of a uniform irradiation after the fall of the meteorite. However, we found a ratio of activity "Mn/" Mn equal to 0.12 ± 0.12. Thus it seems, that the Granès meteorite has been irradiated less strongly during the few days preceding its fall.

*** THE END **

REFERENCES

- [1].- D. NORDEMANN ET J. TOBAILEM. Comptes-Rendus, p. 3581 of T.259, 1964.
- [2].- J. TOBAILEM ET D. NORDEMANN.- Geochimica and Cosmochimica Acta (print)
- [3].- M. HONDA ET D. LAL. Nucl. Physics, 51, p. 363, 1964.
- [4].- G. V. S. RAYUDU. Canadian J. of Chemistry (in print).

Contract No.NAS-5-3760 Consultants & Designers, Inc. Arlington, Virginia Translated by ANDRE L. BRICHANT on 24 September 1965

DISTRIBUTION

GODDARD SPACE FLIGHT CENTER				NASA HQS		OTHER CENTERS		
100 110 400 600 610 611	CLARK, TOWNSE STROUD BOURDEAU PIEPER MEREDITH McDONALD	ND	SS SG	NEWELL, N MITCHELL SCHARDT DUBIN ROMAN SMITH	AUGLE	SONET LIBRA Lar C	RY [3]	
	GUSS KNIFTEN ABRAHAM BOLDT		SL	LIDDEL BRUNK BRYSON FELLOWS		160 213 185	ADAMSON KATZOFF DAVIS WEATHERWAX	[2]
612	HEPPNER NESS			HIPSHER HOROWITZ		JP L	WDATHDAWAA	[4]
613 614	KUPPERIAN LINDSAY WHITE	[3]	SM	FOSTER ALLENBY GILL		NEWBU EIMER SNYDE	!	
615	BAUER GOLDBERG STONE		RR RTR	BADGLEY KURZWEG NEILL		SPEED CONEL KOVAC	en En	
640	HESS O'KEEFE	[3]	ATSS	ROBBINS	[3]	LOOMI UCL		
630 252 256	GI for SS LIBRARY FREAS	[5]	WX	SWEET		COLEM		